

## ■ ARZNEIMITTELPRÜFUNG

Das fluktuierende Krankheitsgeschehen kann den Wirksamkeitsnachweis einer Therapie deutlich erschweren

# Klinische Forschung bei Autoimmunerkrankungen – berücksichtigen wir dabei die „Störfaktoren“ ausreichend?

*Dr. Ines Schöndorf, MBA, SAPERCO-GmbH, Buseck*

## Quellen

[1] Arem H, Moore SC, Patel A et al. Leisure time physical activity and mortality: a detailed pooled analysis of the dose-response relationship. *JAMA Intern Med* 2015; 175:959-967.

[2] Sveaas SH, Smedslund G, Hagen KB, Dagfinrud H. Effect of cardiorespiratory and strength exercises on disease activity in patients with inflammatory rheumatic diseases: a systematic review and meta-analysis. *Br J Sports Med* 2017; 51:1065-1072.

[3] Barry A, Cronin O, Ryan AM et al. Impact of Exercise on Innate Immunity in Multiple Sclerosis Progression and Symptomatology. *Front Physiol* 2016; 7:194.

[4] Canning KL. Physical activity and multiple sklerosis. McMaster University, Hamilton, Ontario; 2017.

[5] Xin J, Feinstein DL, Hejna MJ, Lorens SA, McGuire SO. Beneficial effects of blueberries in experimental autoimmune encephalomyelitis. *J Agric Food Chem* 2012; 60:5743-5748.

[6] Lerner A, Jeremias P, Matthias T. The World Incidence and Prevalence of Autoimmune Diseases is increasing. *Int J Celiac dis* 2015; 3:151-155.

[7] Gioxari A, Kaliora AC, Marantidou F, Panagiotakos DP. Intake of omega-3 polyunsaturated fatty acids in patients with rheumatoid arthritis: A systematic review and meta-analysis. *Nutrition* 2018; 45:114-124.

[8] Petersson S, Philippou E, Rodomar C, Nikiphorou E. The Mediterranean diet, fish oil supplements and Rheumatoid arthritis outcomes: evidence from clinical trials. *Autoimmun Rev* 2018; 17:1105-1114.

[9] Hadgkiss EJ, Jelinek GA, Weiland TJ, Pereira NG, Marck CH, van der Meer DM. The association of diet with quality of life, disability, and relapse rate in an international sample of people with multiple sclerosis. *Nutr Neurosci* 2015; 18:125-136.

[10] Katz S, I. The Role of Diet in Multiple Sclerosis: Mechanistic Connections and Current Evidence. *Curr Nutr Rep* 2018; 7:150-160.

[11] Thompson JM, Li T, Park MK, Qureshi AA, Cho E. Estimated serum vitamin D status, vitamin D intake, and risk of incident alopecia areata among US women. *Arch Dermatol Res* 2016; 308:671-676.

[12] Shapira Y, Agmon-Levin N, Shoenfeld Y. Defining and analyzing geoepidemiology and human autoimmunity. *J Autoimmun* 2010; 34:J168-J177.

[13] Santos-Antunes J, Nunes AC, Lopes S, Macedo G. The Relevance of Vitamin D and Antinuclear Antibodies in Patients with Inflammatory Bowel Disease Under Anti-TNF Treatment: A Prospective Study. *Inflamm Bowel Dis* 2016; 22:1101-1106.

[14] Wang S, Wu Y, Zuo Z, Zhao Y, Wang K. The effect of vitamin D supplementation on thyroid autoantibody levels in the treatment of autoimmune thyroiditis: a systematic review and a meta-analysis. *Endocrine* 2018; 59:499-505.

[15] Li J, Chen N, Wang D, Zhang J, Gong X. Efficacy of vitamin D in treatment of inflammatory bowel disease: A meta-analysis. *Medicine (Baltimore)* 2018; 97:e12662.

[16] Soubrier M, Lambert C, Combe B et al. A randomised, double-blind, placebo-controlled study assessing the efficacy of high doses of vitamin D on functional disability in patients with rheumatoid arthritis. *Clin Exp Rheumatol* 2018; 36:1056-1060.

[17] De Filippo C, Di PM, Ramazzotti M et al. Diet, Environments, and Gut Microbiota. A Preliminary Investigation in Children Living in Rural and Urban Burkina Faso and Italy. *Front Microbiol* 2017;8:1979.

[18] Singh A, Sarangi AN, Goel A et al. Effect of administration of a probiotic preparation on gut microbiota and immune response in healthy women in India: an open-label, single-arm pilot study. *BMC Gastroenterol* 2018; 18:85.

[19] Garg V, Lipka S, Rizvon K, Singh J, Rashid S, Mustacchia P. Diffuse nodular lymphoid hyperplasia of intestine in selective IgG 2 subclass deficiency, autoimmune thyroiditis, and autoimmune hemolytic anemia: case report and literature review. *J Gastrointest Liver Dis* 2012; 21:431-434.

[20] Dargahi N, Johnson J, Donkor O, Vasiljevic T, Apostolopoulos V. Immunomodulatory effects of probiotics: Can they be used to treat allergies and autoimmune diseases? *Maturitas* 2019; 119:25-38.

[21] Rinaldi E, Consonni A, Guidesi E, Elli M, Mantegazza R, Baggi F. Gut microbiota and probiotics: novel immune system modulators in myasthenia gravis? *Ann N Y Acad Sci* 2018; 1413:49-58.

[22] Schorpion A, Kolasinski SL. Can Probiotic Supplements Improve Outcomes in Rheumatoid Arthritis? *Curr Rheumatol Rep* 2017;19:73.

[23] Forbes A, Escher J, Hebuterne X et al. ESPEN guideline: Clinical nutrition in inflammatory bowel disease. *Clin Nutr* 2017; 36:321-347.

[24] Gratton SM, Herro AM, Feuer WJ, Lam BL. Cigarette Smoking and Activities of Daily Living in Ocular Myasthenia Gravis. *J Neuroophthalmol* 2016; 36:37-40.

[25] Roelsgaard IK, Thomsen T, Ostergaard M et al. The effect of an intensive smoking cessation intervention on disease activity in patients with rheumatoid arthritis: study protocol for a randomised controlled trial. *Trials* 2017; 18:570.

[26] Fisher MC, Hochberg MC, El-Taha M, Kremer JM, Peng C, Greenberg JD. Smoking, smoking cessation, and disease activity in a large cohort of patients with rheumatoid arthritis. *J Rheumatol* 2012; 39:904-909.

[27] Andersson ML, Bergman S, Soderlin MK. The Effect of Stopping Smoking on Disease Activity in Rheumatoid Arthritis (RA). Data from BARFOT, a Multicenter Study of Early RA. *Open Rheumatol J* 2012; 6:303-309.

[28] Zhang J, Wei W, Wang CM. Effects of psychological interventions for patients with systemic lupus erythematosus: a systematic review and meta-analysis. *Lupus* 2012; 21:1077-1087.

[29] Schneider-Schaulies J, Beyersdorf N. CD4+ Foxp3+ regulatory T cell-mediated immunomodulation by anti-depressants inhibiting acid sphingomyelinase. *Biol Chem* 2018; 399:1175-1182.

[30] Faissner S, Mishra M, Kaushik DK et al. Systematic screening of generic drugs for progressive multiple sclerosis identifies clomipramine as a promising therapeutic. *Nat Commun* 2017; 8:1990.

[31] Koopman FA, van Maanen MA, Vervoordeldonk MJ, Tak PP. Balancing the autonomic nervous system to reduce inflammation in rheumatoid arthritis. *J Intern Med* 2017; 282:64-75.

[32] Rausch Osthoff AK, Niedermann K, Braun J et al. 2018 EULAR recommendations for physical activity in people with inflammatory arthritis and osteoarthritis. *Ann Rheum Dis* 2018; 77:1251-1260.

[33] Gopalakrishnan V, Spencer CN, Nezi L et al. Gut microbiome modulates response to anti-PD-1 immunotherapy in melanoma patients. *Science* 2018; 359:97-103.

[34] Goubet AG, Daillere R, Routy B, Derosa L, Roberti M, Zitvogel L. The impact of the intestinal microbiota in therapeutic responses against cancer. *C R Biol* 2018; 341:284-289.

[35] Deutsche Rheuma-Liga Bundesverband e.V. aktiv-hoch-r, Das Bewegungsprogram für mehr Lebensqualität. URL: [www.aktiv-hoch-r.de/index.php/aktiv-hoch-r-die-idee](http://www.aktiv-hoch-r.de/index.php/aktiv-hoch-r-die-idee) (letzter Aufruf: 18.01.2019).

ENDE