

■ ARZNEIMITTELPRÜFUNG

Statistik – Folge 3 |

Schlüsselwörter: Primärer Endpunkt, Sekundärer Endpunkt, kombinierter Endpunkt

Primäre Endpunkte und sekundäre Endpunkte

Andreas Ziegler *^a, Katja Krockenberger^b

^a StatSol, Lübeck ^b AMEDON GmbH, Lübeck

* Prof. Dr. Andreas Ziegler, StatSol, Moenring 2, 23560 Lübeck, Tel.: 0451 / 98960279, Fax: 0451 / 98960279, E-Mail: ziegler@statsol.de

Quellen

- [1] Dias J, Brealey S, Choudhary S, et al. Scaphoid Waist Internal Fixation for Fractures Trial (SWIFFT) protocol: a pragmatic multi-centre randomised controlled trial of cast treatment versus surgical fixation for the treatment of bi-cortical, minimally displaced fractures of the scaphoid waist in adults. *BMC Musculoskelet Disord* 2016; 17:248.
- [2] Doehn C, Krockenberger K, Zumbé J, et al. Comparison of white-light, photodynamic diagnosis and narrow-band imaging for the detection of non-muscle invasive bladder cancer: results from a randomized multicenter diagnostic phase-III study. *European Urology Supplements* 2014; 13:e23.
- [3] Huque MF. On some statistical considerations in testing for multiple endpoints in clinical trials. *Science, Statistics and Decisions The Right Questions and the Best Answers*; 2004; Washington D.C.
- [4] Krummenauer F, Al-Nawas B, Baulig C. Primäre klinische Endpunkte: es sollte nur einen geben ... *Zeitschrift für Zahnärztliche Implantologie* 2008; 24:215-217.
- [5] Lengsfeld M, König IR, Schmelter J, et al. Passive rotary dynamic sitting at the workplace by office-workers with lumbar pain - a randomised multi-centre study. *Spine Journal* 2007; 7:531-540; discussion 540.
- [6] Mueller-Godeffroy E, Vonthein R, Ludwig-Seibold C, et al. Psychosocial benefits of insulin pump therapy in children with diabetes type 1 and their families - the pumpkin multicenter randomized controlled trial. *Pediatr Diabetes* 2018; 19:1471-1480.
- [7] Müller-Godeffroy E, Treichel S, Wagner VM, et al. Investigation of quality of life and family burden issues during insulin pump therapy in children with Type 1 diabetes mellitus — a large-scale multicentre pilot study. *Diabetic Medicine* 2009; 26:493-501.

- [8] Nelson PR. Primary and secondary endpoints. In: Itani KMF, Reda DJ, Hrsg. Clinical Trials Design in Operative and Non Operative Invasive Procedures. Cham: Springer; 2017:11-20.
- [9] O'Neill RT. Secondary endpoints cannot be validly analyzed if the primary endpoint does not demonstrate clear statistical significance. Controlled Clinical Trials 1997; 18:550-556; discussion 561-557.
- [10] Pocock SJ, Stone GW. The primary outcome fails — what next? New England Journal of Medicine 2016; 375:861-870.
- [11] Pocock SJ, Stone GW. The primary outcome is positive — is that good enough? New England Journal of Medicine 2016; 375:971-979.
- [12] Schmidt C, Öner A, Mann M, et al. A novel integrated care concept (NICC) versus standard care in the treatment of chronic cardiovascular diseases: protocol for the randomized controlled trial CardioCare MV. Trials 2018; 19:120.
- [13] Ziegler A, Krockenberger K. Patientenrelevante Endpunkte versus Surrogatendpunkte: Klassische Beispiele aus der Literatur. PM QM 2018; 20:142-146.
- [14] Zura R, Xiong Z, Einhorn T, et al. Epidemiology of fracture nonunion in 18 human bones. JAMA Surgery 2016; 151:e162775.

ENDE